

Homogeneity, Flatness and “Large” Extra Dimensions

Glenn D. Starkman¹, Dejan Stojkovic¹ and Mark Trodden²

¹ *Department of Physics
Case Western Reserve University
Cleveland, OH 44106-7079, USA*

² *Department of Physics
Syracuse University
Syracuse, NY, 13244-1130 USA*

We consider a model in which the universe is the direct product of a (3+1)-dimensional Friedmann, Robertson-Walker (FRW) space and a compact hyperbolic manifold (CHM). Standard Model fields are confined to a point in the CHM (i.e. to a brane). In such a space, the decay of massive Kaluza-Klein modes leads to the injection of any initial bulk entropy into the observable (FRW) universe. Both Kolmogoro-Sinai mixing due to the non-integrability of flows on CHMs and the large statistical averaging inherent in the collapse of the initial entropy onto the brane smooth out any initial inhomogeneities in the distribution of matter and of 3-curvature on any slice of constant 3-position. If, as we assume, the initial densities and curvatures in each fundamental correlation volume are drawn from some universal underlying distributions independent of location within the space, then these smoothing mechanisms effectively reduce the density and curvature inhomogeneities projected onto the FRW. This smoothing is sufficient to account for the current homogeneity and flatness of the universe. The fundamental scale of physics can be $\lesssim 1\text{TeV}$. All relevant mass and length scales can have natural values in fundamental units. All large dimensionless numbers, such as the entropy of the universe, are understood as consequences of the topology of spacetime which is not explained. No model for the origin of structure is proffered.

PACS:12.10.-g, 11.10.Kk,
11.25.M,04.50.+h

SU-GP-01/5-1

The notion [1–4] that space has more than three space and one time dimensions, and thus that our 3+1 dimensional universe is merely a subspace (“the brane”) on which ordinary Standard Model (SM) fields are confined inside a higher dimensional space (“the bulk”) in which gravitational degrees of freedom propagate, offers new perspectives for cosmology [5,6]. A simple realization is a space-time which factorizes into an ordinary dynamical (3+1)-dimensional Friedmann-Robertson-Walker universe $\mathcal{M}_{FRW}$ and a static extra-dimensional spatial manifold, $\mathcal{M}_{\text{extra}}$; so that $\mathcal{M} = \mathcal{M}_{FRW} \times \mathcal{M}_{\text{extra}}$. The brane on which standard model fields are localized is taken to be stationary at a point in $\mathcal{M}_{\text{extra}}$. Previously, we showed [5] that if $\mathcal{M}_{\text{extra}}$ has hyperbolic, rather than flat, geometry, then the dynamics of the very early universe alleviate some standard cosmological problems. In this letter we argue that the homogeneity and local flatness of the universe are consequences of a broad class of initial conditions for such a universe, assuming that $\mathcal{M}_{\text{extra}}$ is sufficiently topologically complex and that the fundamental laws of physics are universal.

The principal motivation for large extra dimensions was to solve the gauge hierarchy problem. For an extra dimensional manifold of volume V_{extra} , the force due to massless gravitons will, by Gauss’s law, follow an inverse square law with an effective coupling of

$$M_{\text{pl}}^{-2} = M_F^{-(d+2)} V_{\text{extra}}^{-1} \quad (1)$$

where M_F is the fundamental scale of gravity. The graviton in these theories is a mode which is homogeneous on $\mathcal{M}_{\text{extra}}$. Contributions to the gravitational potential due to excitations of the extra-dimensional geometry (Kaluza-Klein (KK) modes) are Yukawa-like and important only at distances short compared to the inverse mass of the associated KK mode.

For $d = 2$ and 3, most compact manifolds admit only a hyperbolic homogeneous geometry. CHMs are obtained from their universal covering space H^d , by “modding-out” by a (freely-acting) discrete subgroup Γ of the isometry group $SL(d, 1)$ of H^d . If the extra dimensions comprise a CHM [7] then all experimental constraints are met by $M_F \simeq TeV$ and a maximum spatial extent of the CHM of only $\mathcal{O}(30)M_F^{-1}$.

An important property of CHM’s is the dependence of their volume on linear size: $V \sim b_c^d e^\beta$ for a given curvature radius b_c , with βb_c proportional to the maximum linear dimension of the manifold. Crucially, for a homogeneous geometry, e^β is a topological invariant of the manifold; in $d = 2$, it is proportional to the Euler characteristic. In $d = 2$ and 3, there is a countable infinity of CHMs with volumes distributed approximately uniformly from a finite minimum value to infinity (in units of the curvature scale b_c). In large volume CHM’s there is generically a gap in the spectrum of the Laplacian between the zero mode (the graviton) and the next lowest

mode. A theorem due to Sarnak in $d = 2$ and a conjecture due to Brooks in $d \geq 3$ state (approximately) that for large volumes characteristically $m_{\text{gap}} = \mathcal{O}(b_c^{-1})$. The natural value for b_c is $\mathcal{O}(M_F^{-1})$. One therefore expects

$$M_{KK} \gtrsim M_F. \quad (2)$$

The solution to the gauge hierarchy problem in this model is therefore at least technically natural, corresponding to the combination of a large, but topological, value of e^β and the natural value $b_c M_f = \mathcal{O}(1)$.

Aside from its attractive particle-physics features, this construction has important cosmological implications. In the balance of this letter we show how the large scale homogeneity and local flatness of our universe emerge essentially automatically from relatively generic local initial conditions and a specific choice for the topology of spacetime, and hence for e^β .

We imagine that the universe appears, in that its geometry can first be treated classically, at a time t_1 . We assume that any initial evolution of the geometry of $\mathcal{M}_{\text{extra}}$ is brief and ignorable. This assumption is likely justified if, as we assume, the mechanism stabilizing the radius of curvature of $\mathcal{M}_{\text{extra}}$ yields $b_c(t_1) = \mathcal{O}(M_F^{-1})$. The volume of $\mathcal{M}_{\text{extra}}$ is given by (1) with $e^\beta = \mathcal{O}(M_{Pl}^2/M_F^2)$. We do not explain this large value of e^β , but note that it is a topological invariant of $\mathcal{M}_{\text{extra}}$.

We assume further that $\mathcal{M}_{FRW}$ is expanding for $t \geq t_1$ with massive KK modes dominating the energy density of the universe. We assume that these KK modes have initial number density of $g_1 T_1^{3+d}$ (g_1 here being some slowly changing multiplicity factor, and $T_1 \simeq M_F$ some effective temperature), with energy density $g_1 T_1^{3+d} M_{KK}$.

As $\mathcal{M}_{FRW}$ expands, the KK modes are out of thermal equilibrium and eventually decay, ultimately into light modes – gravitons and standard model particles. We show below that in this process most of the entropy in the universe moves from the bulk to the brane. We assume for simplicity that the KK decay is instantaneous, occurring at $t_2 \equiv \tau_{KK} = M_{Pl}^2/(g_{KK} M_{KK}^3)$ (g_{KK} quantifies the multiplicity of KK decay channels). With this approximation, we may equate the total energy content of the comoving 3-volume before and after the decay:

$$e^\beta b_1^d a_1^3 g_1 T_1^{3+d} M_{KK} = g_2 T_2^4 a_2^3 \quad (3)$$

where $a_{1,2}$ are the FRW scale factors at time $t_{1,2}$. Assuming that the universe was effectively matter dominated from $t_1 = [8\pi g_1 T_1^{4+d}/(3M_F^{2+d})]^{-1/2}$ to t_2 , i.e. $(a_2/a_1) = (t_2/t_1)^{2/3}$ we obtain

$$T_2 = (3/(8\pi))^{1/4} g_{KK}^{1/2} g_2^{-1/4} M_{Pl}^{-1/2} M_{KK}^{7/4} T_1^{-1/4}. \quad (4)$$

If $T_2 \gtrsim 1\text{MeV}$, which can easily be realized, nucleosynthesis is unaffected. For example with $d = 3$, $g_1 \simeq g_2 \simeq g_{KK} \sim 100$, $M_{KK} = 3M_F$, $M_F = 5\text{TeV}$,

we find $T_2 > 1\text{MeV}$. For a consistent cosmology, baryogenesis probably must occur during or prior to the decay of the KK modes.

The decay of the KK modes leads to the usual radiation dominated era, from t_2 to t_3 , followed by a matter dominated era, from t_3 to today.

We next calculate the level of residual inhomogeneity we expect in our universe today. We assume that at t_1 , when the initial distribution of KK modes is set, there exists some correlation scale $\xi \simeq M_F^{-1}$, below which fluctuations in all quantities are highly correlated, but above which all fluctuations are completely uncorrelated. We assume further that on this scale there are $\mathcal{O}(1)$ fluctuations about a well defined mean density $\bar{\rho} = \mathcal{O}(M_F^{4+d})$. Most importantly, we assume the universality of the underlying distribution from which the ρ in each correlation volume is drawn.

In momentum space, these fluctuations correspond to a gas of massive KK modes populating the bulk. As $\mathcal{M}_{FRW}$ expands, the gas cools, and the KK modes decay via a cascade into massless modes (i.e. $m \ll M_F$). However, the massive KK modes cannot decay purely into gravitons (or gravitinos) because both KK modes and gravitons are eigenmodes of the generalized wave operator for spin-2 particles on the CHM and are therefore orthogonal on the CHM. Furthermore, since the graviton is the zero-mode of this operator on the CHM, the product of the wavefunctions of any number of gravitons will still be homogeneous. The decay rate for a massive KK state to pure gravitons will involve an overlap integral between the KK wavefunction and the multi-graviton wavefunction and will therefore vanish at lowest order in perturbation theory. This can also be understood in terms of a residual discrete isometry of the CHM, inherited from H^d , under which the KK modes carry non-zero charges while gravitons are scalars. (The need to prevent KK decay to multi-gravitons may constrain deviations from factorizability of $\mathcal{M}$ which break the discrete isometry.)

The brane, however, breaks the residual symmetry. KK modes can decay into Standard model particles which are confined to the brane, and thus are not eigenmodes of the same operator. The coupling of KK modes to Standard Model modes is typically unsuppressed, compared to the coupling between KK modes, except for the wave-function overlap suppression. The KK modes therefore preferentially decay to Standard Model particles propagating on the brane, or to these plus gravitons. Since there are many light non-gravitational modes on the brane, and only one massless gravitational mode in the bulk, we expect most decays of KK modes to deposit their energy and entropy in SM fields on the brane.

Entropy injection into the brane is a powerful homogenizing processes in this cosmology. Consider a primordial fluctuation $\delta\rho/\rho(\xi, t_1)$. So long as the underlying distribution of initial perturbations has sufficiently compact support, the central limit theorem ensures that the mag-

nitude of this fluctuation at some length scale λ at some time t is suppressed by a huge number $\mathcal{O}(\sqrt{n})$, where n is the number of appropriately redshifted initial correlation volumes (ξ^{3+d}) which project into the 3-volume λ^3 :

$$n \simeq \frac{e^\beta b_c^d \lambda^3}{\xi^{3+d} (a(t)/a_1)^3}. \quad (5)$$

The growth of density fluctuations in time partially balances this statistical averaging. Primordial fluctuations grow as $t^{2/3}$ in the matter dominated era while in the radiation dominated era are frozen [8]. Thus, the density fluctuations at some late time ($t > t_3$) at the scale λ are related to the initial ones by:

$$\frac{\delta\rho}{\rho}(\lambda, t) \simeq \frac{\xi^{(3+d)/2} (a(t)/a_1)^{3/2}}{e^{\beta/2} b_c^{d/2} \lambda^{3/2}} \left(\frac{t_2}{t_1}\right)^{2/3} \left(\frac{t}{t_3}\right)^{2/3} \frac{\delta\rho}{\rho}(\xi, t_1)$$

At the time of last scattering t_{ls} , for the fluctuations at the apparent horizon scale, $\lambda = \lambda_{ls}$, we have

$$\frac{\delta\rho}{\rho}(\lambda_{ls}, t_{ls}) \simeq 2.6 \times g_1^{5/6} g_{KK}^{-11/12} g_2^{-3/8} M_{Pl}^{19/12} \xi^{(d+3)/2} \times T_1^{(20d+71)/24} T_3 M_F^{-(d+2)/3} M_{KK}^{-19/8} T_4^{-5/2} \lambda_{ls}^{-3/2} \frac{\delta\rho}{\rho}(\xi, t_1) \quad (6)$$

Here $T_3 = 5.5\Omega_0 h^2 \text{eV}$ is the temperature at the beginning of the matter dominated era, $T_4 = 0.3\text{eV}$ is the temperature of the CMB at last scattering and we take $\lambda_{ls} \sim 3 \times t_{ls}$, with $t_{ls} = 180000(\Omega_0 h^2)^{-1/2} \text{yrs}$. We assume $\Omega_0 = 1$, $h = 0.7$. It is easy to see that the homogeneity of the universe ($\delta\rho/\rho(3t_{ls}) \lesssim 10^{-5}$) can be achieved for reasonable values of parameters. For example, to keep $T_2 > 1\text{MeV}$ we use the values of the relevant parameters given below eq. (4) for which we find $\delta\rho/\rho \sim 4 \cdot 10^{-19}$ (assuming $\frac{\delta\rho}{\rho}(\xi, t_1)$ of order one). Thus homogeneity is achieved without fine tuning of the initial parameters.

Since the power spectrum of fluctuations behaves as $\lambda^{-3/2}$, the scale at which fluctuations become large (of order one) is safely small — of order $4 \times 10^9 \text{m}$.

It is of interest to note that even prior to the homogenization that occurs due to the decay of the KK modes and the injection of the associated entropy into the brane, an additional homogenization mechanism is at work — mixing due to the non-integrability of flows in the CHM, i.e. chaos. CHMs are the paradigmatic arena for chaotic dynamics. Ignoring the effects of gravitational feedback, these dynamics cause gradients in the energy-density to be reduced as $e^{-\kappa d}$, where κ is the Kolmogorov-Sinai (K-S) entropy [9] of the flow, and d is the distance the flow travels. It has been shown [10] that $\kappa \simeq V_{CHM}^{(-1/d)} \log 2$. Using $d \simeq \tau_{KK} \simeq M_{Pl}^2/M_{KK}^3$ and $M_{KK} \simeq M_F$, we find

$$\kappa d \simeq (M_{Pl}/M_F)^{2(1-1/d)} \log 2 / (M_F b_c)$$

This is a huge factor, which suggests that by the time of KK decay, at any given point in $\mathcal{M}_{FRW}$, the density is

incredibly homogeneous across $\mathcal{M}_{extra}$. Note that these flows are effective at eliminating only those gradients of the density which are in the extra-dimensional directions — ordinary gradients are mostly unaffected by this effect, unless $\mathcal{M}_{FRW}$ is also a CHM. Nevertheless, this mechanism assures that the full $\sqrt{n}$ suppression of density fluctuations is realized. (This homogenizing effect of flows has been pointed out in a cosmological context, but with ordinary three-space as the CHM, by other authors looking for alternatives to inflation [11], or at least alternative solutions to the homogeneity problem [12].)

What then of the flatness problem — the fact that the universe today is so very close to flat, with curvature radius $a_c \simeq \frac{3000h^{-1} Mpc}{\sqrt{1-\Omega_0}}$? We must separate this problem into two pieces:

a) The Global Flatness Problem: Is the manifold of three-dimensional space-time one with a topology that admits a flat homogeneous metric, or one with a topology that admits a curved (spherical or hyperbolic) homogeneous metric, or neither? We do not solve this problem, but rather assume that the global topology of space time is consistent with Euclidean geometry and a trivial topology (infinite extent in all directions). This is at least a technically natural assumption, i.e. it is topological invariant unchangeable by cosmological dynamics absent the formation of a singularity.

b) The Local Flatness Problem: Given that the global topology is what it is, why have fluctuations in the local curvature on cosmological scales not caused either the recollapse of the visible universe, or its runaway expansion — i.e. why does the universe look so flat?

In the context of this scenario, the flatness problem is just the homogeneity problem applied to gravity — the effective three-curvature in a particular domain in the FRW dimensions is obtained by averaging over many independent domains in the bulk. The curvature can be either positive or negative on a given domain of radius ξ . If the distribution around zero curvature is symmetric, then by the central limit theorem, the mean 3-curvature today will be given by:

$$\frac{1}{\bar{a}_4^2} = \frac{1}{\sqrt{n}} \frac{1}{a_4^2} = \frac{1}{\sqrt{n}} \frac{1}{a_1^2} \left(\frac{a_1}{a_2}\right)^2 \left(\frac{a_2}{a_3}\right)^2 \left(\frac{a_3}{a_4}\right)^2 \quad (7)$$

where n is calculated at the scale of the visible universe, i.e. the Hubble distance cH_0^{-1} . Expressed in fundamental units, the initial energy density of the universe is

$$G_{4+d}\rho_1 = \frac{g_1}{M_F^{d+2}} T_1^{4+d}, \quad (8)$$

where G_{4+d} is the fundamental (4+d dimensional) gravitational constant and g_1 is the density of states. The appropriately redshifted energy density today is

$$G_{4+d}\rho_4 = \frac{g_1}{M_F^{d+2}} T_1^{4+d} \left(\frac{a_1}{a_2}\right)^3 \left(\frac{a_2}{a_3}\right)^4 \left(\frac{a_3}{a_4}\right)^3, \quad (9)$$

where we have used correspondingly redshifted powers in the matter and radiation dominated eras.

The flatness problem is the smallness of the ratio between $G_{4+d}\rho_4$ and $1/\bar{a}_4^2$; but now

$$\frac{G_{4+d}\rho_4}{1/\bar{a}_4^2} \simeq \lambda^{3/2} M_{Pl}^{-7/12} M_F^{-(2d+4)/3} M_{KK}^{-9/8} T_1^{(4d+37)/24} T_3 \times T_4^{5/2} a_1^2 \xi^{-(d+3)/2} g_{KK}^{-1/12} g_1^{1/6} g_2^{7/8} \quad (10)$$

T_3 is, as before, the temperature at the beginning of the matter dominated era, $T_4 = 3 \cdot 10^{-4} \text{eV}$ is the temperature today while $\lambda = cH_0^{-1} = 9.78h^{-1}10^9 \text{lyrs}$. This ratio is much greater than unity for any reasonable values of parameters. To keep $T_2 \gtrsim 1 \text{MeV}$, we use the values of the relevant parameters as given below (4) for which $T_2 > 1 \text{MeV}$ giving 10^8 . Thus, the curvature term today in Einstein's equation is negligible with respect to the energy density term which achieves flatness without fine tuning of the initial parameters.

The Kolmogorov-Sinai smoothing will play the same role as in the homogeneity problem.

The net effect of these mechanisms is to solve the local aspect of the entropy problem. The entropy within the horizon at some earlier time (say CMBR) was much smaller than the entropy within the horizon today, thus implying that the horizon today consists of many causally disconnected regions, and yet the universe appears uniform. In the context of our homogenization mechanism (and its assumptions about the topology of $\mathcal{M}_{\text{extra}}$), this is no longer a problem. We should note that, since we assumed global Euclidean geometry and trivial topology of the 3-space, the global entropy problem (the existence of the huge dimensionless number of 10^{88} , as discussed in [13]) is equivalent to the global flatness problem — why the universe is so big (infinite in extent). As we stated earlier, we do not solve this problem.

We have thus shown that, under certain assumptions, the homogeneity and local flatness of the visible universe is a generic feature of a universe with large extra dimensions and factorizable geometry. To do so we have had to assume that the underlying distribution functions from which density and curvature are drawn are homogeneous. Additionally, the scenario places certain demands on the topology of space-time — it must be factorizable $\mathcal{M} = \mathcal{M}_{FRW} \times \mathcal{M}_{\text{extra}}$, with $\mathcal{M}_{FRW}$ homeomorphic to $R \times E^3$, and the topology of $\mathcal{M}_{\text{extra}}$, as quantified by e^β , sufficiently complex. Moreover, it is apparent that the topology of $\mathcal{M}_{\text{extra}}$ must not vary, or at least not by a lot, from point to point in $\mathcal{M}_{FRW}$. We make no attempt here to quantify the limits on how much the topology may vary, for it clearly depends on whether the only effect will be a change in the volume, and hence the value of the effective gravitational coupling G_N . In that case the limits are likely quite weak since we know only that $\Delta G_N/G_N \lesssim 0.1$ over the relevant length and time scales. However, if other constants of nature, such

as fermion masses, or α_{EM} are very sensitive to changes in the topology, then more restrictive limits may apply.

We have not asserted any solution for the origin of structure — though we may hope that fluctuations of the brane might be the source — nor have we addressed the issue of baryogenesis — though we note that the decay of the KK modes is out of equilibrium and could be baryon-number and CP violating. Further, we acknowledge that there are difficulties for proton stability, though these have been previously discussed in the context of the canonical scenario [2].

In summary, while we do not claim at present that this is a complete alternative view of a universe with neither fine-tunings nor inflation, we have demonstrated some interesting cosmological consequences of large extra dimensions, warranting further detailed study.

This work was supported by NSF grant PHY-0094122 (MT) and by the DOE (GDS and DS).

-
- [1] I. Antoniadis, Phys. Lett. **B246**, 377 (1990); J. Lykken Phys. Rev. **D54**, 3693 (1996).
 - [2] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B429**, 263 (1998); I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B436**, 257 (1998).
 - [3] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999); *ibid* 4690 (1999); J. Lykken and L. Randall, hep-th/9908076; N. Arkani-Hamed, S. Dimopoulos, G. Dvali, N. Kaloper, Phys. Rev. Lett. **84** 586 (2000).
 - [4] I. Antoniadis and K. Benakli, Phys. Lett. **B326**, 69 (1994); K. Dienes, E. Dudas, and T. Gherghetta, Phys. Lett. **B436**, 55 (1998), Nucl. Phys. **B537**, 47 (1999).
 - [5] G. D. Starkman, D. Stojkovic and M. Trodden, Phys. Rev. D **63**, 103511 (2001)
 - [6] J. Khoury, B. A. Ovrut, P. J. Steinhardt, N. Turok, hep-th/0103239
 - [7] N. Kaloper, J. March-Russell, G.D. Starkman and M. Trodden, Phys. Rev. Lett. **85**, 928 (2000).
 - [8] J. A. Peacock, *Cosmological Physics*, Cambridge University Press, Cambridge (1999).
 - [9] A.N. Kolmogorov, Dokl. Akad. Nauk. SSSR **98**, 527 (1954); Ya.G. Sinai, *ibid.*, **124**, 768 (1959).
 - [10] J.R. Gott, M.N.R.A.S. 193, 153 (1980); C.N. Lockhart, B.Misra & I.Prigogine, Phys. Rev. **D25**, 921 (1982).
 - [11] O. Heckmann & E. Schucking, in *Handbuch der Physik*, ed. S. Flugge (Springer, Berlin, 1959), Vol. 53, p.515; G.F. Ellis, Gen. Rel. Grav. **2**, 7 (1971); H.V. Fagundes, Phys. Rev. Lett. **51**, 517 (1983); D. J. H. Chung, K. Freese, Phys. Rev. **D62** 063513 (2000).
 - [12] N. J. Cornish, D. N. Spergel, G. D. Starkman, Phys. Rev. Lett. **77** 215 (1996)
 - [13] A.D. Linde, *Particle Physics and Cosmology*, section 1.5, (Harwood, Chur, Switzerland, 1990).